



## Examination of State-Trait Anxiety Levels of Smoking Students Preparing for University Exam During the Earthquake

Depremde üniversite sınavına hazırlanan sigara kullanan öğrencilerin durumluk-sürekli kaygı düzeylerinin incelenmesi

 Şeyda Öztuna<sup>1</sup>, Feray Bucak<sup>2</sup>

<sup>1</sup>Balıkesir Atatürk Şehir Hastanesi,

<sup>2</sup>Harran University Faculty of Health Sciences Nursing Department

### ABSTRACT

**Purpose:** This study was conducted to examine the state-trait anxiety levels of students who were preparing for the university exam and were smokers during the earthquake.

**Method:** The study was conducted with 130 people who were preparing for university and smokers who experienced the earthquake in Elazığ Sivrice, the epicenter of which was on January 24, 2020 at 20:55 local time. Data were collected from individuals using the "Individual Information Form" and "State-Trait Anxiety Inventory (STAI I-II)". The obtained data were analyzed with the SPSS 25 program.

**Results:** It was determined that the state anxiety levels of male subjects and the trait anxiety levels of female subjects included in the study were high. Those who had no idea about the effects of smoking on health had significantly higher trait anxiety levels than others. It was determined that there was a positive and low-level significant relationship between state anxiety and trait anxiety. It was found that the effect of the state anxiety level on the trait anxiety level was significant.

**Conclusion:** The study concluded that the level of state-trait anxiety is high in individuals who smoke and have earthquake experience and are preparing for university. It is thought that organizing training programs and prioritizing rehabilitation studies can be effective in preventing and/or reducing anxiety that may develop in individuals after an earthquake and developing effective intervention methods to cope with anxiety.

**Keywords:** Earthquake, State Anxiety, Smoking, Stress, Trait Anxiety.

### ÖZET

**Amaç:** Bu çalışma deprem olgusunda üniversite sınavına hazırlanan ve sigara kullanan öğrencilerin durumluk-sürekli kaygı düzeylerinin incelenmesi amacıyla yapılmıştır.

**Gereç ve Yöntem:** Çalışma 24 Ocak 2020 tarihinde yerel saatle 20.55'te merkez üssü Elazığ Sivrice depremini yaşamış üniversiteye hazırlanan ve sigara kullanan çalışmaya katılmayı kabul eden 130 kişi ile yürütülmüştür. Bireylerden "Birey Bilgi Formu" ve "Durumluk-Sürekli Kaygı Ölçeği (STAI I-II)" kullanılarak veriler toplanmıştır. Elde edilen veriler SPSS 25 programı ile analiz edilmiştir.

**Bulgular:** Çalışmaya dahil edilen erkek olguların durumluk kaygı durumları ve kadın olguların sürekli kaygı durumlarının yüksek olduğu saptanmıştır. Sigaranın sağlık üzerine etkileri hakkında fikri olmayanların diğerlerine göre sürekli kaygı düzeyleri anlamlı şekilde yüksek çıkmıştır. Durumluk kaygı ile sürekli kaygı arasında pozitif yönde düşük düzeyde anlamlı bir ilişki olduğu tespit edilmiştir. Durumluk kaygı düzeyinin sürekli kaygı düzeyine etkisinin anlamlı olduğu bulunmuştur.

**Sonuç:** Çalışma deprem yaşantısı olan sigara kullanan üniversiteye hazırlanan bireylerde durumluk-sürekli kaygı düzeyinin yüksek olduğu sonucuna varılmıştır. Bireylerde deprem sonrasında gelişebilecek kaygının önlenmesi ve/veya azaltılması ve kaygı ile baş edebilmede etkin müdahale yöntemlerinin geliştirilebilmesi amacıyla eğitim programlarının düzenlenmesinin ve rehabilitasyon çalışmalarına öncelik verilmesinin etkili olabileceği düşünülmektedir.

**Anahtar Kelimeler:** Deprem, Durumluk Kaygı, Sigara, Stres, Sürekli Kaygı.

Corresponding Author: Şeyda Öztuna, e-mail: seydaoztuna@gmail.com

Received: 13.10.2024, Accepted: 9.12.2024, Published Online: 15.03.2025

How to cite: Öztuna, Ş., & Bucak, F. (2025). Examination of State-Trait Anxiety Levels of Smoking Students Preparing for University Exam During the Earthquake. *Anatolian Journal of Mental Health*. 2(1):28-48.

<https://doi.org/10.5281/zenodo.14991413>

## INTRODUCTION

The World Health Organization (WHO) defines disaster as “an event that disrupts normal conditions of existence and causes suffering at a level that exceeds the adaptive capacity of the affected society” (WHO, 2022). Disasters have been an important problem for humanity throughout history. Disasters that occur in some parts of the world can cause serious losses at national and international levels and have dramatic effects on people (Taş and Çakır, 2022). Disasters, which are a universal problem, are examined in two sections as natural and human disasters. Natural disasters are the result of natural causes such as hurricanes, earthquakes, tsunamis and tropical cyclones (Makwana, 2019). Earthquakes are events that occur suddenly and cause destruction, injuries and loss of life in homes (Cred, 2020). Earthquakes are the first in the list of natural disasters (Yardım, 2023). The World Health Organization reported that earthquakes between 1998 and 2017 caused approximately 750,000 deaths and more than 125 million people were affected by earthquakes (Igarashi et al., 2021). In Turkey, there were 269 earthquakes that caused loss of life or damage between 1900 and 2023 (Presidency of the Republic of Turkey, Strategy and Budget Presidency, 2023). The Elazığ earthquake with a magnitude of 6.8 occurred on January 24, 2020 at 20:55 local time in the Sivrice district of Elazığ, and 1185 aftershocks were recorded until February 8, 2020 after this main earthquake. 41 people lost their lives and 1632 people were seriously injured in the main earthquake (Özmen et al., 2023). Earthquake phenomena can cause psychological problems in individuals (Nakajima, 2012). Individuals who lose their loved ones and experience economic problems after a major traumatic event such as an earthquake also have an increased risk of experiencing psychological problems such as nightmares, sleep problems, post-traumatic stress disorder, and dissociative reactions (Johannesson et al., 2011; Bıçakçı and Okumuş, 2023). Therefore, individuals may experience feelings such as shock, surprise, anger, helplessness, powerlessness, loss of confidence, loss of control, fear of death, and anxiety (Yıldız and Akkoyun, 2023). The Elazığ Sivrice earthquake caused great material and spiritual destruction, and many problems including the situations listed above were experienced and recorded. In a study, children and adolescents were asked to write an essay covering descriptions of the earthquake after the Tohoku earthquake and the tsunami that occurred immediately after. The essays written by the students were interpreted by the researchers as the appearance of tsunami waves creating anxiety in children and adolescents. Students' themed compositions such as "weakness, nausea and a funny feeling" were also expressed as physical reactions triggered by fear and anxiety (Hayashi et al., 2012). Another problem

encountered in adolescents who experienced an earthquake is smoking and addiction. In a study examining the drug consumption of adolescents affected by the earthquake in the city of Bam in Iran, it was stated that drug use increased after the earthquake (Sharifian et al., 2019).

### **Reasons for Teenagers Smoking**

It is stated that smoking is an important health problem that threatens human health (Zaher et al., 2004). It is estimated that the number of cigarette users in the world will reach 1.7 billion in 2025 (WHO, 2004). Due to the easy accessibility of cigarettes, their legality, and the fact that they pave the way for addictions to other substances such as heroin, alcohol, and marijuana, it can appear as substance addiction (Yakar et al., 2020). In addition to its addictive effect, smoking is the most important preventable cause of mortality and morbidity in our country. WHO reported that 25% of adults worldwide smoke and approximately 80% of those who smoke live in countries including our country (WHO, 2015). According to TUIK 2016 data in Turkey, smoking is reported to be 40.1% in men over the age of 15 and 13.3% in women. It is also estimated that these rates will continue to increase rapidly (TUIK, 2016). It is known that individuals experience smoking during adolescence due to the following situations (Kasatura, 1998; Kassel et al., 2003):

- Thinking that they cannot cope with stress when they quit smoking
- Thinking that they will eat too much when they quit smoking
- Being able to leave negative moods such as fear, anger, and anxiety and switch to a calming, pleasurable and satisfying positive mood
- Taking those who smoke as an example in their environment
- Envyng their friends
- Disaster situations such as earthquakes
- Peer pressure
- Escaping anxiety
- Not being excluded from the social environment
- The idea that one time will not be a problem
- Curiosity
- Wanting to prove that they are grown up
- Growing up in a family that talks about smoking as a stress reliever and a factor that helps to focus
- Thinking that smoking can help them

## **Types of Anxiety**

Anxiety is defined as a feeling of uneasiness about a subjective situation that a person is experiencing in the moment and whose future occurrence is uncertain or will not occur. Individuals can use defense mechanisms by ignoring and rejecting situations that cause them pain (Erskine, 2018). Anxiety is divided into two: state and trait anxiety (Takhdat et al., 2024).

**State Anxiety:** State anxiety is defined as a temporary anxiety reaction given in situations that may cause anxiety (Leal et al., 2017). It can also be expressed as the individual's psychological reactions, provided that the intensity of occurrence is not frequent and continuous. State anxiety can vary from event to event. Examples include birth anxiety, childbearing anxiety, or daily life anxiety (Cruise, et al., 1985).

State anxiety can result from the person's perception of the situation they are in as a threat. This situation creates an unpleasant emotional state in the person. This emotional state can be perceived, understood, and felt.

**Trait Anxiety:** Trait anxiety is the increasing frequency and intensity of reactions to situations that cause stress or that the individual perceives as dangerous. Individuals with high levels of trait anxiety experience stressful or anxiety-provoking events more intensely than those with low levels. Trait anxiety has the following characteristics (Spielberger and Reheiser, 2009):

- It continues for a long time in the individual
- It is continuous
- The duration and intensity of anxiety may vary from person to person
- The psychological discomfort that the individual has may affect the level of trait anxiety
- Individuals' psychological differences may affect the evaluation process of events perceived as dangerous and threatening.

## **Effects of Anxiety on the Individual**

Anxiety is one of the most common psychological problems in the world (Rapee et al., 2023). It is mostly desired and expected for anxiety to be experienced mildly and normally. When there is no anxiety to a certain extent, the person may lack motivation. However, excessive anxiety can cause emotional harm to the individual (Öz, 2017).

It is stated that 301 million people from all age groups have anxiety disorders and the rate of anxiety during adolescence is 12.9% (WHO, 2022). In a study conducted with 917

high school students in Turkey, it was determined that 75% of the participants had anxiety disorders (Özakar Akça et al., 2018). According to a study conducted in the USA in 2012, 34.1% of high school students had anxiety disorder symptoms, while in 2018, 44% of the participants had anxiety disorder symptoms (Parodi et al., 2022).

The anxiety and expectation that emerges in children and adolescents who experience a major earthquake can cause negative reactions such as anxiety. Studies have shown that children and adolescents exposed to natural disasters have high anxiety levels after the event and even for many years (Lai et al., 2015; Shi et al., 2016).

It has been observed that anxiety in adolescents can cause different psychological problems in the future (Tang et al., 2020). It has been stated that anxiety disorder in adolescence is associated with psychological problems such as depression (Melton et al., 2016) and some technology addictions (Li et al., 2019; Yang et al., 2019). Therefore, the symptoms of anxiety disorders can continue for a long time in the person's life (Lau, 2022).

Earthquakes can cause persistent anxiety in individuals and societies (Hayes, 2011). Individuals with high trait anxiety levels have weak cognitive flexibility. These individuals are sensitive to stimuli that may cause anxiety and have difficulty responding harmoniously (Wang et al., 2018). They tend to focus on negativity against stimuli that may cause anxiety. They show relatively high physiological responses to anxious situations (Knowles and Olatunji, 2020). It can be said that anxiety causes significant negative effects on individuals in cognitive, emotional and behavioral dimensions. It is thought that current studies are needed in this area since the desired level of success has not been achieved despite the fight against smoking in Turkey. Based on the gap in the literature on the subject, this study aims to evaluate the state-trait anxiety states of university students who are smokers and who were exposed to the Elazığ Sivrice earthquake on January 24, 2020 at 20:55 local time. It is anticipated that the data obtained from this study, which examines the anxiety states of the students, will be useful in the development of effective stress coping methods due to the fact that Turkey is located in the earthquake zone and smoking is increasing among young people.

The hypothesis of this study is stated below:

**H1:** State-trait anxiety levels of students who smoke while preparing for the university exam may be affected during the earthquake.

## MATERIALS AND METHODS

### Purpose and Type of Research

This research is a descriptive and correlational research methodology in quantitative research methodology. The state-trait anxiety levels of smoking university students who were exposed to the Elazığ Sivrice earthquake at 20:55 local time on January 24, 2020 were evaluated.

### Population and Sample of the Research

The universe of the study consisted of smokers preparing for university who experienced the earthquake with the epicenter in Elazığ Sivrice at 20:55 local time on January 24, 2020; the sample consisted of 130 people who agreed to participate in the study.

### Data Collection Tool

The data of the study was collected with the “Individual Information Form and State-Trait Anxiety Inventory (STAI I-II)”. Before the application, the participants were interviewed face to face and the purpose of the study was explained. The data of the study was obtained using the questionnaire form and scale after obtaining consent from the individuals who agreed to participate in the study. The data was collected face to face and online method (online form) considering the working conditions in line with the demands of the individuals.

**Individual Information Form:** The Individual Information Form is a form prepared by researchers and is used to obtain personal information about the participants. This form includes information on smoking status, number of cigarettes smoked daily, reason for starting smoking, presence of smokers in the family, awareness of the effects of smoking on health, education, employment and socioeconomic status of the participants' parents, and the number of times they have taken the university entrance exam.

**State-Trait Anxiety Scale:** The State-Trait Anxiety Scale, developed by Spielberger et al. (1970), was adapted to Turkish by Öner and Le Compte (1983). According to this scale, if more than three statements are not answered, the completed form is considered invalid and is not scored. Scoring in the State Anxiety Scale is 1=Never, 2=A Little, 3=A Lot, and 4=Completely; scoring in the Trait Anxiety Scale is 1=Almost never, 2=Sometimes, 3=A Lot, and 4=Almost always. There are 40 questions in total in the State-Trait Anxiety Scale. The scores obtained from both scales theoretically vary between 20 and 80. A high score indicates a high level of anxiety, a low score indicates a low level of anxiety. The internal reliability total (Cronbach's alpha) values of the Trait Anxiety Scale vary between 0.83-0.87, and for

state anxiety behavior it is between 0.94-0.96 (Spielberger et al., 1970; Oner and Compte, 1985). With these improvements, Cronbach's alpha numbers change to 0.81 for State Anxiety and 0.68 for Trait Anxiety.

### **Statistical Analysis**

The data of the study were analyzed with the Statistical for Social Science for Windows 25 (SPSS, Version 25; Armonk, NY: IBM Corp. 2017) program. In the normality distribution tests, it was seen that the Skewness and Kurtosis values were less than 1 and the data were found to be normally distributed. In the analysis of the data, descriptive statistics such as number, percentage, mean, standard deviation, minimum, maximum, independent sample T test, One-way ANOVA test, Pearson correlation test and simple regression test were used. The statistical significance level was accepted as  $p < 0.05$ .

### **Ethics Approval**

This study was conducted in accordance with the decision of Balıkesir Atatürk City Hospital Non-Interventional Clinical Research Ethics Committee dated 29.02.2024 and numbered 2024/01/1 and in accordance with the Helsinki protocol.

## **RESULTS**

Table 1 compares the sociodemographic characteristics of the participants and the effects of these characteristics on their anxiety levels. It was determined that 58.5% of the participants were female; 73.1% of their mothers were illiterate; 45.4% were taking the exam for the second time; 96.2% of their mothers were unemployed; 13.8% had more income than expenses and 41.5% took the exam from the equal weight area. The state anxiety levels of the participants were found to be significantly higher in males ( $p = 0.008$ ). Trait anxiety was found to be statistically significantly higher in females and in those whose fathers had a high school or higher education ( $p < 0.05$ ). No significant relationship was found between other sociodemographic characteristics and anxiety levels ( $p > 0.05$ ).

**Table 1. Comparison of State-Trait Anxiety Levels of Participants According to Their Sociodemographic Characteristics**

Sociodemographic Characteristics		n	%	State Anxiety $\bar{x} \pm SD$	Constant Anxiety $\bar{x} \pm SD$
<b>Gender</b>	Female	76	58.5	40.32±5.21	48.11±6.60
	Male	54	41.5	43.03±6.12	43.83±6.56
				<b>p=0.008</b> t=-2.714	<b>p=0.001</b> t=3.657
<b>Mother's Education Status</b>	Illiterate	95	73.1	41.31±5.96	45.86±7.12
	Literate	11	8.5	41.18±6.61	45.81±4.23
	Primary school graduate	18	13.8	41.94±4.89	48.50±7.59
	Secondary school graduate	6	4.6	42.66±3.32	48.33±6.89
				p=0.925 F=0.156	p=0.430 F=0.927
<b>Father's Education Status</b>	Illiterate	16	12.3	42.31±6.73	44.75±10.36
	Literate	18	13.8	42.00±4.91	44.44±4.23
	Primary school graduate	66	50.8	40.86±5.37	45.80±6.21
	Secondary school graduate	22	16.9	40.45±6.11	48.68±6.27
	High school and above	8	6.2	46.12±6.15	51.75±7.47
				p=0.125 F=1.840	<b>p=0.036</b> F=2.649
<b>YGS Entry Number</b>	First time entering	41	31.5	40.14±4.57	46.90±6.20
	Second time entering	59	45.4	42.20±6.12	46.05±7.83
	More than twice entering	30	23.1	41.76±6.25	46.13±5.89
				p=0.201 F=1.625	p=0.819 F=0.199
<b>Does Mother Work?</b>	Yes	5	3.8	43.20±3.27	50.00±5.43
	No	125	96.2	41.38±5.81	46.19±6.92
				p=0.490 t=0.692	p=0.227 t=1.214
<b>Does Father Work?</b>	Yes	92	70.8	40.97±5.87	46.04±6.69
	No	38	29.2	42.62±5.39	47.08±7.48
				p=0.144 t=-1.471	p=0.443 t=-0.769
<b>Family Income Status</b>	Income-expense equal	56	43.1	41.55±5.63	45.10±6.50
	Income-expense more	18	13.8	41.77±5.57	46.94±8.61
	Income-expense less	56	43.1	41.25±5.99	47.37±6.59
				p=0.931 F=0.071	p=0.204 F=1.611
<b>Parents' Attitude</b>	Democratic	38	29.2	40.81±6.13	46.15±5.95
	Authoritarian	23	17.7	39.82±5.82	45.34±5.01
	Protective	60	46.2	42.46±5.60	47.13±7.62
	Indifferent	9	6.9	41.55±4.00	44.33±9.57
				p=0.244 F=1.408	p=0.244 F=0.682
<b>Exam Area</b>	Numerical	25	19.2	41.44±4.42	45.56±6.29
	Verbal	51	39.2	42.33±6.68	47.35±7.95
	Equal Weight	54	41.5	40.62±5.27	45.74±6.03
				p=0.318 F=1.157	p=0.404 F=0.914

(p= .005; p<.05)



Table 2 examines the effects of participants' attitudes towards smoking on state-trait anxiety levels. It was determined that 86.2% of the participants smoked; half of the smokers smoked eleven or more cigarettes per day; 44.4% started smoking due to the influence of friends; 38.6% said their families did not smoke; and 45.4% said smoking caused lung diseases and respiratory distress. It was determined that there was no statistically significant relationship between attitudes towards smoking and state anxiety scores ( $p>0.05$ ). Although the state anxiety levels of those whose parents smoked were higher than the other groups, it was not statistically significant. It was found that those who had no idea about the effects of smoking on health had significantly higher trait anxiety levels than the others ( $p=0.017$ ).

**Table 2. Comparison of the Effects of Participants' Attitudes Towards Smoking on State-Trait Anxiety Levels**

Attitudes Towards Smoking			State Anxiety	Constant Anxiety	
	n	%	$\bar{x} \pm SD$	$\bar{x} \pm SD$	
Smoking	Yes	18	13.8	42.88±7.48	40.22±8.62
	No	112	86.2	41.22±5.42	47.32±6.06
			$p=0.255$ $t=1.143$	$p=0.001$ $t=-4.325$	
Number of Cigarettes Smoked Daily	10 pieces per day and below	9	50.0	42.11±7.80	37.44±9.13
	11 pieces per day and above	9	50.0	43.66±7.53	43.00±7.56
Reasons for Starting Smoking	Peer influence	8	44.4	44.62±8.03	38.75±12.39
	Other (boredom etc.)	10	55.6	41.50±7.12	41.40±4.19
			$p=0.395$ $t=0.874$	$p=0.534$ $t=0.049$	
Smokers in the Family	Only father	48	36.9	42.41±5.82	45.97±7.26
	Only mother	5	3.8	40.00±4.94	43.00±9.19
	Mother and father	5	3.8	43.60±6.22	51.80±4.32
	Sibling	22	16.9	40.54±7.20	47.00±8.03
	No one uses	50	38.6	40.86±4.95	46.18±5.81
			$p=0.491$ $F=0.859$	$p=0.325$ $F=1.174$	
What are the Health Effects of Smoking?	Causes cancer	18	13.8	42.11±4.12	47.11±5.01
	Causes lung diseases and respiratory distress	59	45.4	40.94±6.25	44.79±7.12
	Causes heart disease	17	13.1	41.17±4.74	45.11±7.90
	No idea	36	27.7	42.22±6.11	49.25±6.10
			$p=0.720$ $F=0.447$	$p=0.017$ $F=3.507$	

( $p= .005$ ;  $p<.05$ )

The mean scores of the state anxiety and trait anxiety scales and the correlation analysis between them are presented in Table 3. The mean state anxiety score was found to be  $41.25 \pm 5.74$  and the mean trait anxiety score was found to be  $46.33 \pm 6.89$ . It is seen that there is a low level positive significant relationship between state anxiety and trait anxiety ( $r=0.179$ ,  $p=0.042$ ).

**Table 3. Correlation between State Anxiety Scale and Trait Anxiety Scale Scores**

	Correlation Matrix		Scale Total Score	
	State Anxiety	Constant Anxiety	Min- Max	$\bar{x} \pm SD$
State Anxiety	1	$r=0.179$ $p=0.042$	26.00-56.00	$41.45 \pm 5.74$
Constant Anxiety	$r=0.179$ $p=0.042$	1	16.00-67.00	$46.33 \pm 6.89$

( $p= .005$ ;  $p<.05$ )

In order to determine the effect of state anxiety level on trait anxiety level, simple linear regression analysis was performed and it was determined that the model established for this purpose was significant ( $F=4.225$ ,  $p=0.042$ ). The  $R^2$  value, which is the explanatory power of the model, was determined as 0.032. According to these results, it can be stated that 0.032% of the trait anxiety variable is explained by the state anxiety variable, which is the independent variable in the model (Table 4).

**Table 4. Regression Analysis of State Anxiety Scale and Trait Anxiety Scale Scores**

Predicted Variable	Predictor Variable	B	SD	$\beta$	t	R	$R^2$	F	Model p
State Anxiety	Constant	34.551	3.395		10.178	0.179	0.032	4.225	<b>0.042</b>
	Constant Anxiety	0.149	0.072	0.179	2.056				

( $p= .005$ ;  $p<.05$ )

## DISCUSSION

In this section, the findings of the research conducted to examine the state-trait anxiety levels of students who were preparing for the university exam and were smoking during the earthquake whose epicenter was Elazığ Sivrice on January 24, 2020 at 20.55 local time were discussed. The effects of sociodemographic variables and smoking on state-trait anxiety were examined. The findings obtained as a result of the analyzes carried out based on the hypotheses of the research were discussed in the light of the studies in the literature.

### **Analysis of Findings Regarding State-Trait Anxiety Levels of Participants According to Their Sociodemographic Characteristics**

It is stated that some demographic factors such as gender, geographical region, residence of parents, occupation of father, family members, occupational status and general satisfaction may affect cigarette consumption (Oktay et al., 2013).

### **Examination of Findings Regarding State-Trait Anxiety Levels According to Gender Factor**

In a study investigating the factors affecting the smoking behaviors of medical and dentistry faculty students, it was determined that being male was among the factors that led students to smoke (Kara et al., 2011). Hoerster (2012) examined the smoking behaviors of students studying at a university and concluded that men smoked more than women. In addition, in this study, it was determined that students who did not live with their families had a higher rate of smoking. In a study investigating anxiety and depression in individuals who experienced an earthquake, it was found that men had higher trait anxiety levels (Miral et al., 1998). A total of 130 people, 76 (58.5) women and 54 (41.5) men, participated in our study conducted to examine the state-trait anxiety levels of students who smoked while preparing for the university entrance exam during the earthquake. It was determined that the state anxiety levels of the participants were significantly higher in men ( $p=0.008$ ). The loss of resources needed to sustain life due to the earthquake, individual differences and the differences in meanings attributed to events may have caused this situation. There are studies reporting that state and/or trait anxiety scores are higher in women than in men (Agun et al., 1993; Dönmez et al., 1996). In different studies, it has been found that female students are more anxious than male students (Keklik, 2011; Deveci et al., 2012; Dursun and Aytaç, 2012; Basco and Olea, 2013; Yılmaz et al., 2014). It is stated that the reason for the prevalence in women is because smoking starts in childhood and increases as the developmental process progresses (Rapee et

al., 2009). Some studies have found that women are at higher risk of experiencing anxiety and depressive disorders (McLean and Anderson, 2009; Parker and Brotchie, 2010; Costello et al., 2011; Schuch et al., 2014). Thus, it has been stated that people may have higher anxiety levels due to feeling exhausted (Otacıoğlu, 2008). Our study data show that trait anxiety is higher in women ( $p < 0.05$ ). It is thought that genetic and biological factors play a role in the emergence of this difference (Zhou et al., 2013). In this study, the high trait anxiety level of women may be due to the effects of biopsychosocial differences, gender roles, hormonal changes being felt more intensely, the belief that women live under relative pressure in society, their inability to use their coping skills effectively, the individual having an anxiety disorder before the earthquake, women being more exposed to the devastating effects of the earthquake and internalizing their anxiety instead of expressing it. There is a need for research that reveals the effect of gender on smoking in individuals who experienced an earthquake.

### **Examination of Findings Regarding State-Trait Anxiety Levels According to Parents' Education Level**

A study conducted to examine the smoking behaviors of university students found that the mother's illiteracy affected the smoking behavior of their children (Özcebe et al., 2014). Yurt Öncel et al. (2011), in their study aiming to determine the risk factors for smoking in university students, determined that the mother's education level had a significant effect on the smoking behavior of children.

Although there are studies indicating that there is no significant relationship between the education level of parents and the anxiety levels of children (Doğan and Çoban, 2009; Tekindal et al., 2010), it is also stated that the trait anxiety level in children decreases as the mother's education level increases (Erdik and Altıparmak, 2012). In our study findings, anxiety levels were found to be statistically significantly higher in those whose fathers' education level was high school and above ( $p < 0.05$ ). This situation may be caused by the relatively high earthquake awareness of the fathers of the participants, and the expectation of an aftershock and the chronic effect of this expectation. In addition, Nolen-Hoeksema and Morrow (1991) stated in their study that pre-earthquake mental illness symptoms are an indicator of post-earthquake PTSD. Therefore, individuals having a chronic or psychiatric illness before may cause anxiety and post-traumatic stress. Since our study did not provide data on individuals' anxiety and post-traumatic stress symptoms before the earthquake, the presence of previous chronic and psychiatric illnesses may have also caused this situation.

### **Examination of Findings Regarding State-Trait Anxiety Levels According to Income Status**

It is stated that income level affects smoking status (Çalışkan, 2015) and that perceiving one's income level as low increases the risk of smoking (Siahpush et al., 2006). Deveci et al. (2010) examined the factors affecting the smoking behavior of students studying at a university and stated that the parents of students who smoke have higher incomes. In some studies investigating the frequency of smoking according to where individuals live, it was found that those with low income levels and living in disadvantaged areas smoke more (Priebe et al., 2009; Nelson et al., 2011).

Our data show that 96.2% of the participants' mothers did not work, 29.2% did not work, and 43.1% had incomes less than expenses. The fact that the socioeconomic status and the state-trait anxiety levels were significantly high in male and female students can be explained by the fact that the students' needs cannot be met or they have difficulty in meeting them, they have to continue their lives in the house where they experienced the earthquake due to financial difficulties, and they cannot use their coping mechanisms effectively because they are relatively more affected by economic losses.

### **Examining the Findings Regarding the Effect of Participants' Attitudes Towards Smoking on Anxiety Levels**

It is stated that individuals may develop automatic negative thinking after disasters such as earthquakes and use smoking as a tool to cope with this situation (Kleinke et al., 1983). There are many studies showing that anxiety is related to smoking (Zvolensky et al., 2003; Breslau et al., 2004; Goodwin et al., 2005). It is reported that university students who smoke have higher trait anxiety scale scores (Comeau et al., 2001; Wiggert et al., 2016). Studies focusing on the effect of smoking on anxiety show that people's emotions return to normal during smoking, but there are deviations from normal in emotions when they do not smoke (Parrott, 1999).

Studies have reported that post-traumatic stress disorder (PTSD) and other psychiatric problems occur at high rates in children and adolescents after an earthquake. After the Northridge earthquake, 28.6% of children developed mild to moderate PTSD (Asarnow et al., 1999); after the Wenchuan earthquake in China, 1.3% of high school students had PTSD symptoms (Zhang et al., 2012); after the earthquake in Haiti in 2010, the prevalence of PTSD and depression in children and adolescents was 36.9% and 46.2% (Cenat and Derivois, 2015); and after the earthquake in Nepal in 2015, 43.3% of adolescents developed probable PTSD

and 38.1% developed probable depression (Sharma and Kar, 2019). In a study conducted with individuals who applied to a Medical Faculty Child Psychiatry Polyclinic after the 1999 Marmara earthquake, it was observed that psychological problems such as communication and elimination disorders were experienced (Berkem and Bildik, 2001).

Our data show that those who had no idea about the effects of smoking on health had significantly higher trait anxiety levels. Depressive moods may have developed in students preparing for university after the earthquake. Therefore, our study findings can be considered as students perceiving smoking as a reward mechanism that reduces trait anxiety and using it as a coping method and experiencing problems in the communication process after the earthquake.

Since no previous research has been conducted on the subject, it is thought that it is important to conduct research to evaluate the state-trait anxiety states of individuals who smoked after the earthquake.

### **CONCLUSION AND RECOMMENDATIONS**

As a result, in our study, where it was determined that some descriptive data differences may have an effect on the state-trait anxiety levels of individuals preparing for university who smoke, we think that preventive activities for smoking may be important. Therefore, according to the findings of our research, we think that conducting different studies and implementing the following measures may be useful in solving smoking:

Within the scope of preventive activities, using a larger sample to determine the state-trait anxiety states of individuals who smoke and have experienced an earthquake with the variables included in our individual information form,

It has been stated that people with alcohol and substance use disorders experience unpleasant emotions such as stress, fear and anger more and tend to use substances to cope with these negative situations (Carpenter et al., 2020). It is stated that interventions to support appropriate coping skills of individuals who have experienced an earthquake should be planned in a way that covers the mental well-being of individuals in line with certain goals,

It is stated that quality sleep, balanced nutrition, avoiding alcohol and substance use, providing physical activity, talking to others about the causes of stress and keeping a diary are effective in reducing stress for individuals who have experienced an earthquake (Babaie et al., 2021; Dakhil et al., 2023). Therefore, health professionals should organize appropriate psychosocial support programs for individuals who have experienced an earthquake to cope

with stress, and the efficiency level of these programs and the feasibility of the targeted situation should be monitored,

Multifaceted, qualitative studies can be conducted in order to offer solutions to smoking. Qualitative studies should examine in detail the effects of different ages, income and education levels, some cultural and social characteristics, and individuals' individual and family situations on determining the factors that cause anxiety and on the use of coping mechanisms for anxiety,

In order to prevent social learning in smoking, appropriate policies should be developed to prevent smoking in places where children are present and to increase smoke-free air spaces,

In some studies conducted after the September 11 events in the United States, it was stated that there was a significant relationship between individuals' television viewing status and anxiety and PTSD symptoms, and that PTSD increased in those who watched these broadcasts (Otto et al., 2007; Bernstein et al., 2007). Therefore, considering that inappropriate news content can affect children and adolescents, parents should check the content their children watch on news channels and social media after the earthquake,

### **Research Statement**

**Ethical Approval:** Balıkesir Atatürk City Hospital Non-Interventional Clinical Research Ethics Committee dated 29.02.2024 and numbered 2024/01/1

**Conflict of Interest:** The authors declare that there is no conflict of interest for the study.

**Financial Support:** This study has received no grants from any funding agency in the public, commercial or social-profit sectors.

### **REFERENCES**

- Agun, K., Saltık, A., Abay, E., Yorulmaz, F., Çağatay, T. ve Tutluoğlu, B. (1993). Üniversite Öğrencilerinde Sınav Stresinin Solunum Fonksiyonlarına Etkisi. *Nöropsikiatri Arşivi*, İstanbul Tıp Fakültesi, İstanbul, 3, 370-375.
- Asarnow, J., Glynn, S., Pynoos, R. S., Nahum, J., Guthrie, D., Cantwell, D. P. ve Franklin, B. (1999). When the earth stops shaking: Earthquake sequelae among children diagnosed for pre-earthquake psychopathology. *Journal of the American Academy of Child & Adolescent Psychiatry*, 38(8), 1016-1023.
- Babaie, J., Naghipour, B. ve Faridaalae, G. (2021). Cardiovascular diseases in natural disasters; a systematic review. *Archives of academic emergency medicine*, 9(1).

- Basco, R. E. ve Olea, M. T. (2013). Correlation between anxiety level and academic performance of BS biology freshmen students. *International Journal of Educational Research and Technology*, 4(1), 97-103.
- Berkem, M. ve Bildik, T. (2001). Depremin Marmara Üniversitesi Tıp Fakültesi Çocuk Psikiyatrisi Polikliniği'ne başvuru profili üzerine etkisi. *Anadolu Psikiyatri Dergisi*, 2(1), 29-35.
- Bıçakcı, A. B. ve Okumuş, F. E. E. (2023). Depremin psikolojik etkileri ve yardım çalışanları. *Avrasya Dosyası*, 14(1), 206-236.
- Breslau, N., Novak, S. P. ve Kessler, R. C. (2004). Daily smoking and the subsequent onset of psychiatric disorders. *Psychological Medicine*, 34(2), 323-333.
- Cénat, J. M. ve Derivois, D. (2015). Long-term outcomes among child and adolescent survivors of the 2010 Haitian earthquake. *Depression and Anxiety*, 32(1), 57-63.
- Comeau, N., Stewart, S. H. ve Loba, P. (2001). The relations of trait anxiety, anxiety sensitivity, and sensation seeking to adolescents' motivations for alcohol, cigarette, and marijuana use. *Addictive Behaviors*, 26(6), 803-825.
- Cred, U. (2020). Centre for Research on the Epidemiology of Disasters. UN Office for Disaster Risk Reduction.
- Çalışkan, Ş. (2015). Üniversite öğrencilerinin sigara kullanımını etkileyen faktörler (Ekonometrik bir yaklaşım). *Uşak Üniversitesi Sosyal Bilimler Dergisi*, 8(2), 23-48.
- Dakhil, ZA, Farhan, HA, Faraz, F., Skuk, MR, Al-Jorani, MS, Rehman, MEU, & Öz, TK (2023). Depremin kardiyovasküler sağlık üzerindeki etkisi: Türkiye ve Suriye'deki yıkıcı depremlerden sonra kardiyovasküler sağlık hizmeti sağlayıcıları ne beklemeli? *Güncel Kardiyoloji Sorunları*, 48(9), 101800.
- Deveci, S. E., Açık, Y., Oguzoncul, A. F. ve Deveci, F. (2010). Prevalence and factors affecting the use of tobacco, alcohol and addictive substance among university students in eastern Turkey. *Southeast Asian J Trop Med Public Health*, 41(4), 996-1007.
- Deveci, S. E., Çalmaz, A. ve Açık, Y. (2012). Doğu Anadolu'da yeni açılan bir üniversitenin öğrencilerinde kaygı düzeylerinin sağlık, sosyal ve demografik faktörler ile ilişkisi. *Dicle Tıp Dergisi*, 39(2), 189-196.
- Doğan, T. ve Çoban, A. E. (2009). Eğitim fakültesi öğrencilerinin öğretmenlik mesleğine yönelik tutumları ile kaygı düzeyleri arasındaki ilişkinin incelenmesi. *Eğitim ve Bilim*, 34(153).



- Dönmez, L., Aktekin, M., Erengin, H., Dinç, G. ve Karaman, T. (1996). Antalya'daki hekimler ve tıp fakültesi öğrencilerinde durumluk ve sürekli kaygı düzeyleri. *Psikiyatri Psikoloji Psikofarmakoloji Dergisi*, 4(4), 268-275.
- Dursun, S. ve Aytaç, S. (2009). Üniversite öğrencileri arasında işsizlik kaygısı. *Uludağ Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 28(1), 71-84.
- Dünya Sağlık Örgütü. (2022). World mental health report: transforming mental health for all.
- Goodman, E. ve Capitman, J. (2000). Ergenlerde depresif semptomlar ve sigara içimi. *Pediatrics*, 106(4), 748-755.
- Gudmundsson, G., Gislason, T., Janson, C., Lindberg, E., Hallin, R., Ulrik, C. S., Brøndum, E., Nieminen, MM., Aine, T. ve Bakke, P. (2005). Risk factors for rehospitalisation in COPD: role of health status, anxiety and depression. *European Respiratory Journal*, 26(3), 414-419.
- Hayes, N. (2011). Psikolojiyi anlamak (F. Şar, Çev.). Optimist Yayın ve Dağıtım.
- Hayashi, K. ve Tomita, N. (2012). Lessons learned from the Great East Japan Earthquake: impact on child and adolescent health. *Asia Pacific Journal of Public Health*, 24(4), 681-688.
- Hoerster, V. (2012). Smoking Behaviour among College Students: A Survey, A Thesis Submitted to the Faculty of Baylor University In Partial Fulfillment of the Requirements for the Honors Program, 1-61,
- Igarashi, Y., Matsumoto, N., Kubo, T., Yamaguchi, M., Nakae, R., Onda, H., Yokobori, Ş., Koido, Y. ve Yokota, H. (2022). Prevalence and characteristics of earthquake-related head injuries: A systematic review. *Disaster Medicine And Public Health Preparedness*, 16(3), 1253-1258.
- Johannesson, E., Simrén, M., Strid, H., Bajor, A. ve Sadik, R. (2011). Physical activity improves symptoms in irritable bowel syndrome: a randomized controlled trial. *Official Journal of the American College of Gastroenterology| ACG*, 106(5), 915-922.
- Kara, S., Baş, F. Y. ve Açıkalın, C. (2011). Sigara içme davranışları ve etkili faktörler: Tıp ve Diş Hekimliği Fakültelerinin ilk ve son sınıf öğrencileri üzerinde çalışma. *Smyrna Tıp Dergisi*, 1(1), 16-21.
- Kasatura, İ. (1998). Gençlik ve Bağımlılık, Evrim Yayınevi ve Bilgisayar San. Tic. Ltd. Şti. İstanbul.

- Keklik, İ. (2011). Relationships of Attachment Styles and Gender, Personal Meaning, Depressiveness and Trait-Anxiety among College Students Üniversite Öğrencilerinin Bağlanma Stilllerinin Cinsiyet, Kişisel Anlamlılık, Depresyona Yatkınlık ve Sürekli Kaygı Düzeyleriyle İlişkisi. *Education*, 36(159).
- Knowles, K. A. ve Olatunji, B. O. (2020). Specificity of trait anxiety in anxiety and depression: Meta-analysis of the State-Trait Anxiety Inventory. *Clinical Psychology Review*, 82, 101928.
- Lai, B. S., Kelley, M. L., Harrison, K. M., Thompson, J. E. ve Self-Brown, S. (2015). Posttraumatic stress, anxiety, and depression symptoms among children after Hurricane Katrina: A latent profile analysis. *Journal of Child And Family Studies*, 24, 1262-1270.
- Lau, WK (2022). Depresyon ve anksiyete semptomlarında dayanıklılığın rolü: Üç dalga çapraz gecikmeli bir çalışma. *Stres ve Sağlık*, 38(4), 804-812.
- Leal, P. C., Goes, T. C., da Silva, L. C. F. ve Teixeira-Silva, F. (2017). Trait vs. state anxiety in different threatening situations. *Trends in Psychiatry And Psychotherapy*, 39(3), 147-157.
- Li, G., Hou, G., Yang, D., Jian, H. ve Wang, W. (2019). Relationship between anxiety, depression, sex, obesity, and internet addiction in Chinese adolescents: A short-term longitudinal study. *Addictive Behaviors*, 90, 421-427.
- Makwana, N. (2019). Disaster and its impact on mental health: A narrative review. *Journal of Family Medicine and Primary Care*, 8(10), 3090-3095.
- Melton, TH, Croarkin, PE, Strawn, JR ve McClintock, SM (2016). Çocuk ve ergenlerde eşlik eden anksiyete ve depresif semptomlar: sistematik bir inceleme ve analiz. *Journal of Psychiatric Practice*, 22(2), 84-98.
- Miral, S., Özcan, Ö., Baykara, A., Yemez, B. ve Tayyar, C. (1998). Dinar depremi sonrası çocuklarda kaygı ve depresyon. *Çocuk ve Gençlik Ruh Sağlığı Dergisi*, 5(1), 16-22.
- Nakajima, Ş. (2012). Deprem ve sonrası psikolojisi. *Okmeydanı Tıp Dergisi*, 28(2), 150-155.
- Nelson, R., Paynter, J. ve Arroll, B. (2011). “Factors Influencing Cigarette access Behaviour Among 14–15-year-olds in New Zealand: A Cross-Sectional Study”, *Journal of Primary Health Care*, 3(2), 114-122.
- Nolen-Hoeksema, S. ve Morrow, J. (1991). A prospective study of depression and posttraumatic stress symptoms after a natural disaster: the 1989 Loma Prieta Earthquake. *Journal of Personality and Social Psychology*, 61(1), 115.

- Oktay, E., Çelik, A. K. ve Akbaba, A. İ. (2013). Examining Demographic Factors Related to Cigarette Smoking among Undergraduate Students at a Turkish University. *International Journal of Higher Education*, 2(2), 175-183.
- Oner N, Le Compte A. Durumluk-Surekli kaygi envanteri el kitabi. Istanbul: Boğaziçi Yayınları; 1985.
- Otto, M. W., Henin, A., Hirshfeld-Becker, D. R., Pollack, M. H., Biederman, J. ve Rosenbaum, J. F. (2007). Posttraumatic stress disorder symptoms following media exposure to tragic events: Impact of 9/11 on children at risk for anxiety disorders. *Journal of Anxiety Disorders*, 21(7), 888-902.
- Öz, S. (2017). Sigara kullanan bireyler ile kullanmayan bireylerin stresle başa çıkma becerileri ve anksiyete düzeylerinin incelenmesi. *Sosyal Bilimler Enstitüsü Dergisi*, 64.
- Akça, S. Ö., Selen, F., Demir, E. ve Demir, T. (2018). Cinsiyet ve yaş farklılıklarının ergenlerin depresyon, anksiyete bozukluğu, kendine zarar verme, psikoz, travma sonrası stres bozukluğu, alkol-uyuşturucu bağımlılığı ve dikkat eksikliği hiperaktivite bozukluğu ile ilişkili sorunlara etkisi. *Dicle Tıp Dergisi*, 45(3), 255-264.
- Özcebe, H., Dogan, B. G., İnal, E., Haznedaroglu, D. ve Bertan, M. (2014). Smoking habits and the related sociodemographic characteristics in university students. *Thoracic Research and Practice*, 15(2), 42.
- Özgüven, İ. E. (1994). *Psikolojik Testler*, Ankara: Yeni Doğu Matbaası.
- Özmen, A., Maraş, M. M., Ayaz, Y. ve Sayın, E. (2023). Assessments of masonry buildings and historical structures during the 2020 Sivrice-Elazığ Earthquake. *Periodica Polytechnica Civil Engineering*, 67(2), 530-544.
- Parodi, K. B., Holt, M. K., Green, J. G., Porche, M. V., Koenig, B. & Xuan, Z. (2022). Time trends and disparities in anxiety among adolescents, 2012–2018. *Social Psychiatry and Psychiatric Epidemiology*, 57(1), 127-137.
- Parrott, A. C. (1999). Does cigarette smoking cause stress?. *American Psychologist*, 54, 817-820.
- Grappasonni, I., Mari, M., Dewey, M., Petrelli, F., & Costa, A. (2009). Posttraumatic stress disorder six months after an earthquake. *Social Psychiatry and Psychiatric Epidemiology*, 44(5), 393.
- Rapee, R. M., Creswell, C., Kendall, P. C., Pine, D. S., & Waters, A. M. (2023). Anxiety disorders in children and adolescents: A summary and overview of the literature. *Behaviour Research and Therapy*, 104376.

- Sharifian, H., Roholamini, M., & Mossavi, M. M. (2019). The Effect Of Earthquake on Youth Tendency to Artificial Narcotics Consumption in Bam Township. *International Journal of Advanced Studies in Humanities and Social Science*, 8(1), 119-124.
- Sharma, A., & Kar, N. (2019). Posttraumatic stress, depression, and coping following the 2015 Nepal earthquake: a study on adolescents. *Disaster Medicine and Public Health Preparedness*, 13(2), 236-242.
- Shi, X., Zhou, Y., & Fan, F. (2016). Longitudinal trajectories and predictors of anxiety symptoms among adolescent survivors exposed to Wenchuan earthquake. *Journal of Adolescence*, 53, 55-63.
- Spielberger CD, Gorsuch RL, Lushene RE, Vagg PR, Jacobs GA. State-trait anxiety inventory. Palo AltoCA: Consulting Psychologists Press; 1970.
- Spielberger, C. D., & Reheiser, E. C. (2009). Assessment of emotions: Anxiety, anger, depression, and curiosity. *Applied Psychology: Health and Well-Being*, 1(3), 271-302.
- T.C. Cumhurbaşkanlığı, Strateji ve Bütçe Başkanlığı, (2023). Kahramanmaraş ve Hatay Depremleri Raporu, Mart, 2023
- Takhdad, K., Rebahi, H., Rooney, D. M., Babram, M. A., Benali, A., Touzani, S., Lamtali, S., & El Adib, A. R. (2024). The impact of brief mindfulness meditation on anxiety, cognitive load, and teamwork in emergency simulation training: A randomized controlled trial. *Nurse Education Today*, 132, 106005.
- Tang, W., Xu, D., & Xu, J. (2020). Impact of earthquake exposure, family adversity and peer problems on anxiety-related emotional disorders in adolescent survivors three years after the Ya'an earthquake. *Journal of Affective Disorders*, 273, 215-222.
- Tas, F., & Cakir, M. (2022). Nurses' knowledge levels and preparedness for disasters: A systematic review. *International Journal of Disaster Risk Reduction*, 80, 103230.
- Tekindal, M., Eryaş, N. ve Tekindal, B. (2010). İlköğretim okullarındaki öğrencilerin sürekli kaygı düzeylerinin çeşitli değişkenlere göre incelenmesi. *Gazi Üniversitesi Endüstriyel Sanatlar Eğitim Fakültesi Dergisi*, 26, 79-93.
- Wang, T., Li, M., Xu, S., Liu, B., Wu, T., Lu, F., Xie, J., Peng, L., & Wang, J. (2019). Relations between trait anxiety and depression: A mediated moderation model. *Journal of Affective Disorders*, 244, 217-222.
- World Health Organization (WHO). "Tobacco free initiative. Building blocks for tobacco control: A Handbook; WHO: Geneva, Switzerland, 2004". [https://www.who.int/tobacco/resources/publications/tobaccocontrol\\_handbook/en/12.10.2019](https://www.who.int/tobacco/resources/publications/tobaccocontrol_handbook/en/12.10.2019).

- World Health Organization (WHO). “WHO report on the global tobacco epidemic, 2015: Raising taxes on tobacco”. [https://apps.who.int/iris/bitstream/handle/10665/178574/9789240694606\\_eng.pdf;jsessionid=07588D7406255D04027A7E97ACE4B01A?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/178574/9789240694606_eng.pdf;jsessionid=07588D7406255D04027A7E97ACE4B01A?sequence=1)/ 12.10.2019.
- Yang, X., Zhou, Z., Liu, Q., & Fan, C. (2019). Mobile phone addiction and adolescents' anxiety and depression: The moderating role of mindfulness. *Journal of Child And Family Studies*, 28, 822-830.
- Yardım, N. (2023). Kahramanmaraş ve Hatay 2023 Depreminden 24 Yıl Önce Marmara Depreminde Gölyaka ve Düzce Çadırkentlerinin Halk Sağlığı Yönünden Değerlendirilmesi. *Afet ve Risk Dergisi*, 6(4), 1368-1387.
- Yıldız, B., & Akkoyun, A. Z. (2023). Deprem sonrası psikiyatrik destek. *İzmir Katip Çelebi Üniversitesi Sağlık Bilimleri Fakültesi Dergisi*, 8(2), 817-820.
- Yılmaz, İ. A., Dursun, S., Güzeler, E. G., ve Pektaş, K. (2014). Üniversite öğrencilerinin kaygı düzeyinin belirlenmesi: bir örnek çalışma. *Ejovoc (Electronic Journal of Vocational Colleges)*, 4(4), 16-26.
- Öncel, S. Y., Gebizlioğlu, Ö. L., ve Alioğlu, F. A. (2011). Risk factors for smoking behavior among university students. *Turkish Journal of Medical Sciences*, 41(6), 1071-1080.
- Zaher, C., Halbert, R., Dubois, R., George, D., & Nonikov, D. (2004). Smoking-related diseases: the importance of COPD. *The International Journal of Tuberculosis and Lung Disease*, 8(12), 1423-1428.
- Zhang, Z., Ran, M. S., Li, Y. H., Ou, G. J., Gong, R. R., Li, R. H., Fan, M., Jiang, Z., & Fang, D. Z. (2012). Prevalence of post-traumatic stress disorder among adolescents after the Wenchuan earthquake in China. *Psychological Medicine*, 42(8), 1687-1693.
- Zhou, X., Kang, L., Sun, X., Song, H., Mao, W., Huang, X., Zhang, Y., & Li, J. (2013). Risk factors of mental illness among adult survivors after the Wenchuan earthquake. *Social Psychiatry and Psychiatric Epidemiology*, 48, 907-915.
- Zvolensky, M. J., Schmidt, N. B., & McCreary, B. T. (2003). The impact of smoking on panic disorder: An initial investigation of a pathoplastic relationship. *Journal of Anxiety Disorders*, 17(4), 447-460.